
Anti-Human/Mouse/Rat Phospho-ERBB2 (Tyr877) Polyclonal Antibody

Polyclonal Antibody

Cat.NO.: PA01600

3th Edition

Description:Protein tyrosine kinase that is part of several cell surface receptor complexes, but that apparently needs a coreceptor for ligand binding. Essential component of a neuregulin-receptor complex, although neuregulins do not interact with it alone. GP30 is a potential ligand for this receptor. Regulates outgrowth and stabilization of peripheral microtubules (MTs). Upon ERBB2 activation, the MEMO1-RHOA-DIAPH1 signaling pathway elicits the phosphorylation and thus the inhibition of GSK3B at cell membrane. This prevents the phosphorylation of APC and CLASP2, allowing its association with the cell membrane. In turn, membrane-bound APC allows the localization of MACF1 to the cell membrane, which is required for microtubule capture and stabilization. In the nucleus is involved in transcriptional regulation. Associates with the 5'-TCAAATTC-3' sequence in the PTGS2/COX-2 promoter and activates its transcription. Implicated in transcriptional activation of CDKN1A; the function involves STAT3 and SRC. Involved in the transcription of rRNA genes by RNA Pol I and enhances protein synthesis and cell growth.

Antigen:Synthesized peptide derived from human Neu around the phosphorylation site of Y877.

Form:

How to use:1.0 ml distilled water will be added to the product

Stability: Lyophilized product, 5 years at 2 – 8°C; Solution, 2 years at –20°C

Dilution:PBS (pH7.4) containing 1% BSA

Application:This antibody can be used for western blotting in concentration of 1?5?g/ml.

Specificity:Expressed in a variety of tumor tissues including primary breast tumors and tumors from small bowel, esophagus, kidney and mouth.