

BHMT2, 1-363aa, Human, His tag, E.coli

Cat.NO.: TP01333

3th Edition

Synonyms:S-methylmethionine-homocysteine S-methyltransferase BHMT2 isoform 1 , betaine--homocysteine S-methyltransferase 2

Description:BHMT2 is involved in the regulation of homocysteine metabolism. Homocysteine is a sulfur-containing amino acid that plays a crucial role in methylation reactions. Transfer of the methyl group from betaine to homocysteine creates methionine, which donates the methyl group to methylate DNA, proteins, lipids, and other intracellular metabolites. BHMT2 is one of two methyl transferases that can catalyze the transfer of the methyl group from betaine to homocysteine. Recombinant human BHMT2 protein, fused to His-tag at N-terminus, was expressed in E.coli.

Form:Liquid. In 20mM Tris-HCl buffer (pH 8.0) containing 0.4M Urea, 10% glycerol

Molecular Weight:42.7 kDa (386aa)

Sequences:

MGSSHHHHHSSGLVPRGSHMGSMAPAGRPGAKKGILERLESGEVVIGDGSFLITLEKRGYVKAGLWTPEAVIEH
PDAVRQLHMEFLRAGSNVMQTFTFSASEDNMESKWEDVNAAACDLAREVAGKGDALVAGGICQTSIYKYQKDEA
RIKKLFRQQLEVFAWKNVDFLIAEYFEHVEEAVWAVEVLKESDRPVAVTCIGPEGMDHITPGECAVRLVKAGASI
VGVNCRFGPDTSLKTMELMKEGLEWAGLKAHLMVQPLGFHAPDCGKEGFVDLPEYPFGLSESRVATRWDIQKYAR
EAYNLGVRYIGGCCGFEPYHIRAIAEELAPERGFPPASEKHGSGWGSGLDMHTKPWIRARARREYWENLLPASGR
PFCPSLSKPDF

Purity:> 95% by HPLC

Concentration:0.5 mg/ml (determined by Bradford assay)

Endotoxin Level:<1.0 EU per 1 ug of protein (determined by LAL method)

Storage:Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.