

**BTD, 44-545aa, Human, His-tag, Baculovirus**

**Cat.NO.: TP01373**

3th Edition

**Synonyms:**Biotinidase isoform 1, BTD, biotinidase

**Description:**BTD, also known as biotinidase isoform 1, is a member of the nitrilase superfamily, which consists of 12 families of nitrilases, amidases, carbamylases, and N-acyltransferases. It catalyzes the hydrolysis of biocytin, the product of biotin-dependent carboxylase degradation, to biotin and lysine. It may have an important regulatory role in chromatin/DNA function. Recombinant human BTD, fused to His-tag at C-terminus, was expressed in insect cell and purified by using conventional chromatography techniques.

**Form:**Liquid. In Phosphate Buffered Saline (pH 7.4) containing 10% glycerol.

**Molecular Weight:**57.8kDa (510aa), 50-70kDa (SDS-PAGE under reducing conditions)

**Sequences:**

```
AHTGEESVADHHEAEYYVAAVYEHPSILSLNPLALISRQEALMLNQNLDIYEQQVMTAAQKDVQIIVFPEDGIHGFN
FTRTSIYPFLDFMPSPQVVRWNPCLPHRFNDTEVLQRLSCMAIRGDMFLVANLGTKEPCHSSDPRCPKDGRYQF
NTNVVFSNNGTLVDRYRKHNLYFEAAFDVPLKVDLITFDTPFAGRFGIFTCFDILFFDPAIRVLRDYKVKHVVPYPTAW
MNQLPLLAIEIQKAFAVAFGINVLAANVHHPVLGMTGSGIHTPLESFWYHDMENPKSHLIIAQVAKNPVGLIGAENA
TGETDPSHSKFLKILSGDPYCEKDAQEVHCDEATKWNVNAPPTFHSEMMYDNFTLVPVWGKEGYLHVCSNGLCC
YLLYERPTLSKELYALGVFDGLHTVHGTYYIQVCALVRCGGLGFDTCGQEITEATGIFEFHLWGNFSTSYIFPLFLTS
GMTLEVPDQLGWENDHYFLRKSRLSSGLVTAALYGRLYERDLEHHHHHH
```

**Purity:**> 95% by HPLC

**Concentration:**0.25mg/ml (determined by Absorbance at 280nm)

**Endotoxin Level:**<1.0 EU per 1 ug of protein (determined by LAL method)

**Storage:**Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.