

CCL18, 22-89aa, Human, His tag, E.coli

Cat.NO.: TP01455

3th Edition

Synonyms:C-C motif chemokine 18, AMAC-1, AMAC1, CKb7, DC-CK1, DCCK1, MIP-4, PARC, SCYA18

Description:CCL18, also known as chemokine (C-C motif) ligand 18, a chemotactic factor that attracts lymphocytes but not monocytes or granulocytes may be involved in B cell migration into B cell follicles in lymph nodes. It attracts naive T lymphocytes toward dendritic cells and activated macrophages in lymph nodes, has chemotactic activity for naive T cells, CD4+ and CD8+ T cells and thus may play a role in both humoral and cell-mediated immunity responses.

Form:Liquid. In 10mM Sodium Citrate pH3.5, 10% Glycerol

Molecular Weight:10.4 kDa (93aa), confirmed by MALDI-TOF

Sequences:

MGSSHHHHHSSGLVPRGSHMGSHMQVGTNKELCCLVYTSWQIPQKFIVDYSETSPQCPCPGVILLTKRGRQICA
DPNKKWVQKYISDLKLNA

Purity:> 95% by HPLC

Concentration:0.25 mg/ml (determined by BCA assay)

Endotoxin Level:<1.0 EU per 1 ug of protein (determined by LAL method)

Storage:Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.