

**CD2, 25-209aa, Human, His tag, E.coli****Cat.NO.: TP01484**

3th Edition

**Synonyms:**T-cell surface antigen CD2, SRBC, T11

**Description:**CD2, also known as T-cell surface antigen T11/Leu-5, is a cell adhesion molecule found on the surface of T cells and natural killer (NK) cells. It interacts with other adhesion molecules, such as lymphocyte function-associated antigen-3 (LFA-3/CD58) in humans, or CD48 in rodents, which are expressed on the surfaces of other cells. In addition to its adhesive properties, CD2 also acts as a co-stimulatory molecule on T and NK cells. CD2 is a specific marker for T cells and NK cells, and can therefore be used in immunohistochemistry to identify the presence of such cells in tissue sections. The great majority of T cell lymphomas and leukaemias also express CD2, making it possible to use the presence of the antigen to distinguish these conditions from B cell neoplasms. Recombinant human CD2 protein, fused to His-tag at N-terminus, was expressed in E.coli.

**Form:**Liquid. 20mM Tris-HCl buffer (pH8.0) containing 10% glycerol 0.4M Urea**Molecular Weight:**23.8 kDa (209aa)**Sequences:**

MGSSHHHHHHSSGLVPRGSHMGSMKEITNALETWGALGQDINLDIPSFQMSDDIDDIKWEKTSKDKKIAQFRKEKE  
TFKEKDTYKLFKNGTLKIKHLKTDDQDIYKVSIDYDTKGKNVLEKIFDLKIQERVSKPKISWTCINTTLTCEVMNGTDPEL  
NLYQDGKHLKLSQRVITHKWTTSLSAKFKCTAGNKVSKESSVEPVSCPEKGLD

**Purity:**> 95% by HPLC**Concentration:**1 mg/ml (determined by Bradford assay)**Endotoxin Level:**<1.0 EU per 1 ug of protein (determined by LAL method)

**Storage:**Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.