

COPS7A, 1-275aa, Human, His tag, E.coli**Cat.NO.: TP01688**

3th Edition

Synonyms:COP9 signalosome complex subunit 7a , CSN7A

Description:COPS7A is a component of the COP9 signalosome complex (CSN), a complex involved in various cellular and developmental processes. The CSN complex is an essential regulator of the ubiquitin (Ubl) conjugation pathway by mediating the deneddylation of the cullin subunits of SCF-type E3 ligase complexes, leading to decrease the Ubl ligase activity of SCF-type complexes such as SCF, CSA or DDB2. The complex is also involved in phosphorylation of p53/TP53, JUN, I-kappa-B-alpha/NFKBIA, ITPK1 and IRF8/ICSBP, possibly via its association with CK2 and PKD kinases. CSN-dependent phosphorylation of TP53 and JUN promotes and protects degradation by the Ubl system, respectively. Recombinant human COPS7A protein, fused to His-tag at N-terminus, was expressed in E.coli.

Form:Liquid. In 20mM Tris-HCl buffer (pH 8.0) containing 0.4M Urea, 10% glycerol

Molecular Weight:32.7 kDa (298aa)

Sequences:

MGSSHHHHHHSSGLVPRGSHMGMSAEVKVTGQNQEQLLLAKSAKGAALATLIHQVLEAPGVVYVFGELLDMPN
VRELAESDFASTFRLLTVFAYGTYADYLAEARNLPPLTEAQKNKLRHLSVVTLAAKVKCIPYAVLLEALALRNVRQLE
DLVIEAVYADVLRGSLDQRNQRLEVDYSIGRDIQRQDLSAIARTLQEWCVGCEVVLSGIEEQVSRANQHKEQQLGL
KQKIESEVANLKKTIKVTAAAAAATSQDPEQHLTELREPAPGTNQRQPSKKASKGKGLRGS AKIWSKSN

Purity:> 95% by HPLC

Concentration:0.5 mg/ml (determined by Bradford assay)

Endotoxin Level:<1.0 EU per 1 ug of protein (determined by LAL method)

Storage:Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.