

## Instruction manual FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

DRG1, 1-367aa, Human, His tag, E.coli

Cat.NO.: TP01922

3th Edition

Synonyms: Developmentally regulated GTP binding protein 1, NEDD3

**Description**:Developmentally-regulated GTP-binding protein 1(DRG1) belong to the GTP1/OBG family. It plays a role in cell proliferation and differentiation, as well as in apoptosis, suggesting a role in tumor formation and metastasis. Expression of the DRG1 was significantly reduced in breast tumor cells, particularly in patients with lymph node or bone metastasis as compared to those with localized breast cancer.

Form:Liquid. In 20mM Tris-HCI buffer (pH 8.0) containing 1mM DTT, 30% glycerol, 1mM EDTA

Molecular Weight: 42.7kDa (387aa), confirmed by MALDI-TOF

## Sequences:

MGSSHHHHHHSSGLVPRGSHMSSTLAKIAEIEAEMARTQKNKATAHHLGLLKARLAKLRRELITPKGGGGGGPGE GFDVAKTGDARIGFVGFPSVGKSTLLSNLAGVYSEVAAYEFTTLTTVPGVIRYKGAKIQLLDLPGIIEGAKDGKGRGR QVIAVARTCNLILIVLDVLKPLGHKKIIENELEGFGIRLNSKPPNIGFKKKDKGGINLTATCPQSELDAETVKSILAEYKIH NADVTLRSDATADDLIDVVEGNRVYIPCIYVLNKIDQISIEELDIIYKVPHCVPISAHHRWNFDDLLEKIWDYLKLVRIYT KPKGQLPDYTSPVVLPYSRTTVEDFCMKIHKNLIKEFKYALVWGLSVKHNPQKVGKDHTLEDEDVIQIVKK

Purity:> 95% by HPLC

Concentration: 0.5mg/ml (determined by Bradford assay)

Endotoxin Level:<1.0 EU per 1 ug of protein (determined by LAL method)

**Storage:**Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.

1/1