
Recombinant Mouse Inducible T-cell Costimulator Protein

Cat.NO.: TP04556

3th Edition

Synonyms:Inducible T-cell costimulator; Activation-inducible lymphocyte immunomediatory molecule; CD28 and CTLA-4-like protein; CCLP; CD28-related protein 1; CRP-1; CD278; Icos; Ailim

Description:Inducible Costimulator(ICOS) is a member of the growing CD28 family of immune costimulatory receptors. Other family members are CD28, CTLA4 and PD1. ICOS shares approximately 39% amino acid similarity with CD 28 and CTLA4. Mouse and human ICOS share approximately 72% amino acid identity. ICOS is expressed on most CD45RO+ cells. ICOS expression is up-regulated within approximately 24-48 hours of activation on Th primed cells. B7-H2, a member of the B7 family of costimulatory ligands, has been identified as the ICOS ligand. The B7-H2/ ICOS interaction appears to play roles in T cell dependent B cell activation and Th differentiation. In addition, ICOS is more potent in the induction of IL-10 production, acytokine important for suppressive function of T regulatory cells.

Form:PBS

Molecular Weight:14.7 kDa

Sequences:Glu21-Leu142

Purity:> 95% by HPLC

Concentration:

Endotoxin Level:<1.0 EU per 1 ug of protein (determined by LAL method)

Storage:Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.