

Recombinant Human Activin Receptor 1A/Activin RI/ALK-2/ACVR1 Protein(C-6His)

Cat.NO.: TP04665

3th Edition

Synonyms:Activin Receptor Type-1; Activin Receptor Type I; ACTR-I; Activin Receptor-Like Kinase 2; ALK-2; Serine/Threonine-Protein Kinase Receptor R1; SKR1; TGF-B Superfamily Receptor Type I; TSR-I; ACVR1; ACVRLK2

Description:Activin receptor type-1, also known as Activin receptor type I, Activin receptor-like kinase 2, Serine/threonine-protein kinase receptor R1, TGF-B superfamily receptor type I, ACVRLK2 and ACVR1, is a single-pass type I membrane protein. ACVR1 is expressed in normal parenchymal cells, endothelial cells, fibroblasts and tumor-derived epithelial cells. ACVR1 belongs to the protein kinase superfamily. Activins signal through a heteromeric complex of receptor serine kinases which include at least two type I (I and IB) and two type II (II and IIB) receptors. These receptors are all transmembrane proteins, composed of a ligand-binding extracellular domain with cysteine-rich region, a transmembrane domain, and a cytoplasmic domain with predicted serine/threonine specificity. Type I receptors are essential for signaling; and type II receptors are required for binding ligands and for expression of type I receptors. Type I and II receptors form a stable complex after ligand binding, resulting in phosphorylation of type I receptors by type II receptors. ACVR1 signals a particular transcriptional response in concert with activin type II receptors.

Form:PBS

Molecular Weight:12.6 kDa

Sequences:Met21-Val124

Purity:> 95% by HPLC

Concentration:

Endotoxin Level:<1.0 EU per 1 ug of protein (determined by LAL method)

Storage:Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.