

## Instruction manual FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

## **Recombinant Mouse CD27 antigen Protein**

Cat.NO.: TP04720

3th Edition

**Synonyms:**CD27 antigen; CD27L receptor; T-cell activation antigen CD27; Tumor necrosis factor receptor superfamily member 7; CD27

**Description:**Mouse CD27 is a single-pass type I membrane protein which is a member of the tumor necrosis factor receptor superfamily. CD27 is required for generation and long-term maintenance of T cell immunity. It binds to ligand CD70, and plays a key role in regulating B-cell activation and immunoglobulin synthesis. CD27 transduces signals that lead to the activation of NF-?B and MAPK8/JNK. Adaptor proteins TRAF2 and TRAF5 have been shown to mediate the signaling process of this receptor. CD27-binding protein (SIVA), a proapoptotic protein, can bind to this receptor and is thought to play an important role in the apoptosis induced by this receptor.

Form:PBS

Molecular Weight: 45.8 kDa

Sequences: Met 1-Arg 182

Purity:> 95% by HPLC

**Concentration:** 

Endotoxin Level:<1.0 EU per 1 ug of protein (determined by LAL method)

**Storage:**Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.

1/1