

**Recombinant Human Leukocyte Ig-Like Receptor B2/LILRB2/ILT4/CD85d Protein(C-6His)**

**Cat.NO.: TP04781**

3th Edition

**Synonyms:**Leukocyte Immunoglobulin-Like Receptor Subfamily B Member 2; LIR-2; Leukocyte Immunoglobulin-Like Receptor 2; CD85 Antigen-Like Family Member D; Immunoglobulin-Like Transcript 4; ILT-4; Monocyte/Macrophage Immunoglobulin-Like Receptor 10; MIR-10; CD85d; LILRB2; ILT4; LIR2; MIR10

**Description:**Members of the immunoglobulin-like transcript (ILT) family are activating and inhibitory immunoreceptors whose genes are located same locus that encodes killer cell Ig-like receptors (KIR). Leukocyte Immunoglobulin-Like Receptor Subfamily B Member 2 (LIR-2) is a type I transmembrane protein. LIR-2 is expressed primarily on monocytes and dendritic cells (DC). Human LIR-2 is produced as a 598 amino acid precursor including a 21 aa signal sequence, a 440 aa extracellular domain (ECD), a 21 aa transmembrane segment, and a 116 aa cytoplasmic domain. LIR-2 binds to Classical MHCI proteins. Ligation of LIR-2 includes Tyr phosphorylation within its cytoplasmic ITIMs, a requirement for association with SHP-1. LIR-2 mediates tolerogenic DC-induced CD4+ T cell energy in vitro and in vivo.

**Form:**PBS

**Molecular Weight:**48.6 kDa

**Sequences:**Gln22-His458

**Purity:**> 95% by HPLC

**Concentration:**

**Endotoxin Level:**<1.0 EU per 1 ug of protein (determined by LAL method)

**Storage:**Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.