

Instruction manual FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

Recombinant Human Death Receptor 3/DR3/TNFRSF25 Protein(C-Fc)

Cat.NO.: TP04935

3th Edition

Synonyms:Tumor necrosis factor receptor superfamily member 25;APO3; DDR3; DR3; TNFRSF12; WSL; WSL1;TNFRSF25;Protein WSL-1;LARD;Protein WSL;Lymphocyte-associated receptor of death;Death receptor 3;Apoptosis-mediating receptor TRAMP;Apoptosis-inducing receptor AIR;Apoptosis-mediating receptor DR3;Apo-3

Description:Tumor necrosis factor receptor superfamily member 25 (TNFRSF25) contains 1 death domain and 4 TNFR-Cys repeats. TNFRSF25 is a cell surface receptor of the tumor necrosis factor receptor superfamily which mediates apoptotic signalling and differentiation, activated by a monogamous ligand, known as TL1A (TNFSF15), which is rapidly upregulated in antigen presenting cells and some endothelial cells following Toll-Like Receptor or Fc receptor activation. This receptor has been shown to signal both through the TRADD adaptor molecule to stimulate NF-kappa B activity or through the FADD adaptor molecule to stimulate caspase activation and regulate cell apoptosis. It may play a role in regulating lymphocyte homeostasis.

Form:PBS

Molecular Weight: 46.3 kDa

Sequences: Gln25-Phe201

Purity:> 95% by HPLC

Concentration:

Endotoxin Level:<1.0 EU per 1 ug of protein (determined by LAL method)

Storage:Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.

1/1