

Instruction manual FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

Recombinant Rat Granulocyte-Macrophage Colony-Stimulating Factor/GM-CSF Protein(C-6His)

Cat.NO.: TP04951

3th Edition

Synonyms:Granulocyte-Macrophage Colony-Stimulating Factor; GM-CSF; Colony-Stimulating Factor; CSF;Molgramostin; Sargramostim; CSF2; GMCSF

Description:Granulocyte-Macrophage Colony Stimulating Factor (GM-CSF) was initially characterized as a growth factorthat can support the in vitro colony formation of granulocyte-macrophage progenitors. It is produced by anumber of different cell types (including activated T cells, B cells, macrophages, mast cells, endothelial cellsand fibroblasts) in response to cytokine of immune and inflammatory stimuli. Besides granulocyte-macrophageprogenitors, GM-CSF is also a growth factor for erythroid, megakaryocyte and eosinophil progenitors. Onmature hematopoietic, monocytes/ macrophages and eosinophils. GM-CSF has a functional role on nonhematopoitic cells. It can induce human endothelial cells to migrate and proliferate. Additionally, GM-CSF canalso stimulate the proliferation of a number of tumor cell lines, including osteogenic sarcoma, carcinoma andadenocarcinoma cell lines.

Form:PBS

Molecular Weight: 15.6 kDa

Sequences: Ala18-Lys144

Purity:> 95% by HPLC

Concentration:

Endotoxin Level:<1.0 EU per 1 ug of protein (determined by LAL method)

Storage:Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.

1/1