

Instruction manual FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

Recombinant Human Ephrin-A1/EFNA1/LERK-1 Protein(C-6His)

Cat.NO.: TP05428

3th Edition

Synonyms:Ephrin-A1; EPH-Related Receptor Tyrosine Kinase Ligand 1; LERK-1; Immediate Early Response Protein B61; Tumor Necrosis Factor Alpha-Induced Protein 4; TNF Alpha-Induced Protein 4; EFNA1; EPLG1; LERK1; TNFAIP4

Description:Ephrin-A1 is a member of the A-type ephrin family of cell surface proteins that function as ligands for the A-type Eph receptor tyrosine kinase family. Ephrin-A1 can be induced by TNF and IL1B. Its expression levels can be down-regulated in primary glioma tissues compared to the normal tissues. The soluble monomeric form is expressed in the glioblastoma multiforme (GBM) and breast cancer cells. Soluble Ephrin-A1 is necessary for the transformation of HeLa and SK-BR3 cells and participates in the relocalization of EPHA2 away from sites of cell-cell contact during transformation. Ephrin-A1 plays an important role in angiogenesis and tumor neovascularization.

Form:PBS

Molecular Weight: 20.4 kDa

Sequences: Asp19-Ser182

Purity:> 95% by HPLC

Concentration:

Endotoxin Level:<1.0 EU per 1 ug of protein (determined by LAL method)

Storage:Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.

1/1