

Recombinant Human Chloride Intracellular Channel Protein 2/CLIC2 Protein(N-6His)

Cat.NO.: TP05712

3th Edition

Synonyms:Chloride Intracellular Channel Protein 2; XAP121; CLIC2

Description:Chloride Intracellular Channel Protein 2 (CLIC2) is a critical component of all living cells; it regulates cellular traffic of Chloride ion and it can be inserted into membranes and form chloride ion channels. Membrane insertion seems to be redox-regulated and may occur only under oxidizing conditions, channel activity depends on the pH. CLIC2 is involved in regulating membrane potential and organic solute transport. CLIC2 modulates the activity of RYR2 and inhibits Calcium influx. CLIC2 can be detected in the adult brain, liver, lung, heart, stomach, spleen and testis. It is expressed in fetal liver and adult skeletal muscle. CLIC2 is a potential candidate for one of many diseases linked to Xq28.

Form:PBS

Molecular Weight:30.5 kDa

Sequences:Met 1-Ser247

Purity:> 95% by HPLC

Concentration:

Endotoxin Level:<1.0 EU per 1 ug of protein (determined by LAL method)

Storage:Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.