

Instruction manual FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

Recombinant Human CFL2 / cofilin 2 / ADF Protein (His tag)

Cat.NO.: TP06256

3th Edition

Synonyms:NEM7

Description:Cofilin 2 (muscle), also known as CFL2, is a member of cofilin family of the actin-binding protein superfamily. Cofilin2 shows significant homology to the other two members: cofilin 1 and DSTN, through its entire sequence, and contains residues conserved among the cofilin family that are responsible for actin-binding. Cofilin 2 (CFL2) is an important regulator of striated myocyte function. Purified cofilin 2 depolymerized actin filaments in a dose- and pH-dependent manner and reduced the apparent viscosity of an actin solution, although they did not cosediment with actin filaments at all. Cofilin2 is not expressed in vegetative cells, but is transiently induced during the aggregation stage of development, whereas cofilin 1 was predominantly expressed in vegetative cells.

Form:PBS

Molecular Weight: 20.4 kDa

Sequences: Ala 2-Leu 166

Purity:> 95% by HPLC

Concentration:

Endotoxin Level:<1.0 EU per 1 ug of protein (determined by LAL method)

Storage:Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.

1/1