
Recombinant Human Contactin 3 / CNTN3 Protein (708 Asp/Asn, His tag)**Cat.NO.: TP07376**

3th Edition

Synonyms:BIG-1;PANG;PCS

Description:Contactins are a subgroup of molecules belonging to the immunoglobulin superfamily that are expressed exclusively in the nervous system. The subgroup consists of six members: Contactin-1, Contactin-2(TAG-1), Contactin-3(BIG-1), BIG-2, Contactin-5(NB-2) and NB-3. Since their identification in the late 1980s, Contactin-1 and Contactin-2 have been studied extensively. Axonal expression and the neurite extension activity of Contactin-1 and Contactin-2 attracted researchers to study the function of these molecules in axon guidance during development. Contactin-1 and Contactin-2 have come to be known as the principal molecules in the function and maintenance of myelinated neurons. In contrast, the function of the other four members of this subgroup remained unknown until recently. Contactin-3, also known as CNTN3 (BIG-1 in rat and PANG in mouse), is a GPI-linked glycoprotein that is expressed on cerebellar Purkinje cells, amygdaloid and thalamic neurons and olfactory granule cells. In the brain, Contactin-3 is expressed in frontal lobe, occipital lobe, cerebellum and amygdala. Contactin-3 contains 4 fibronectin type-III domains and 6 Ig-like C2-type (immunoglobulin-like) domains. Human Contactin-3 shares 92% aa identity with mouse Contactin-3. The exact function of Contactin-3 is unclear. Contactin-3 may mediate cell-cell interaction and may promote neurite outgrowth.

Form:PBS**Molecular Weight:**109.2 kDa**Sequences:**Met 1-Ser1002, 708 Asp/Asn**Purity:**> 95% by HPLC**Concentration:****Endotoxin Level:**<1.0 EU per 1 ug of protein (determined by LAL method)**Storage:**Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.