

Instruction manual FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

Recombinant Human Cochlin / COCH Protein (His tag)

Cat.NO.: TP07551

3th Edition

Synonyms:COCH-5B2;COCH5B2;DFNA9

Description:Cochlin, also known as COCH-5B2 and COCH, is a secreted protein which contains one LCCL domain and two VWFA domains. It is an abundant inner ear protein expressed as multiple isoforms. Its function is also unknown, but it is suspected to be an extracellular matrix component. Cochlin and type II collagen are major constituents of the inner ear extracellular matrix, and Cochlin constitutes 70% of non-collagenous protein in the inner ear, the cochlin isoforms can be classified into three subgroups, p63s, p44s and p40s. The expression of cochlin is highly specific to the inner ear. Eleven missense mutation and one in-frame deletion have been reported in the COCH gene, causing hereditary progressive sensorineural hearing loss and vestibular dysfunction, deafness autosomal dominant type 9 (DFNA9). The co-localization of cochlin and type II collagen in the fibrillar substance in the subepithelial area indicate that cochlin may play a role in the structural homeostasis of the vestibule acting in concert with the fibrillar type II collagen bundles. Defects in COCH may contribute to Meniere disease which is an autosomal dominant disorder characterized by hearing loss associated with episodic vertigo.

Form:PBS

Molecular Weight:59.4 kDa

Sequences: Glu 25-Gln 550

Purity:> 95% by HPLC

Concentration:

Endotoxin Level:<1.0 EU per 1 ug of protein (determined by LAL method)

Storage:Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.

1/1