

---

**Recombinant Human EphA7 / EHK3 Protein (His & GST tag)****Cat.NO.: TP08377**

3th Edition

**Synonyms:**EHK-3;EHK3;EK11;EPA7;HEK11

**Description:**Ephrin type-A receptor 7, also known as EphA7, belongs to the ephrin receptor subfamily of the protein-tyrosine kinase family which 16 known receptors (14 found in mammals) are involved: EPA1, EPA2, EPA3, EPA4, EPA5, EPA6, EPA7, EPA8, EPA9, EPA10, EPB1, EPB2, EPB3, EPB4, EPB5, EPB6. The Eph family of receptor tyrosine kinases (comprising EphA and EphB receptors) has been implicated in synapse formation and the regulation of synaptic function and plasticity<sup>6</sup>. Eph receptor-mediated signaling, which is triggered by ephrins<sup>7</sup>, probably modifies the properties of synapses during synaptic activation and remodeling. Ephrin receptors are components of cell signalling pathways involved in animal growth and development, forming the largest sub-family of receptor tyrosine kinases (RTKs). Ligand-mediated activation of Ephs induce various important downstream effects and Eph receptors have been studied for their potential roles in the development of cancer. Down-regulation of EphA7 secondary to hypermethylation has been reported in colorectal cancer. The expression of EphA7 was reduced in all tested gastric cancer cell lines; however, there is marked variability in expression among gastric carcinoma specimens. EphA7 may have roles in the pathogenesis and development of gastric carcinomas.

**Form:**PBS**Molecular Weight:**75.2 kDa**Sequences:**Gly579-Val998**Purity:**> 95% by HPLC**Concentration:****Endotoxin Level:**<1.0 EU per 1 ug of protein (determined by LAL method)**Storage:**Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.