

Recombinant Human DDR2 Kinase / CD167b Protein (aa 422-855, His & GST tag)

Cat.NO.: TP08447

3th Edition

Synonyms:CD167;MIG20a;NTRKR3;TKT;TYRO10

Description:Discoidin domain receptor 2 (DDR2) or CD167b (cluster of differentiation 167b) is a kind of protein tyrosine kinases associated with cell proliferation and tumor metastasis, and collagen, identified as a ligand for DDR2, up-regulates matrix metalloproteinase 1 (MMP-1) and MMP-2 expression in cellular matrix. DDR2/CD167b was found to recognise the triple-helical region of collagen X as well as the NC1 domain. Binding to the collagenous region was dependent on the triple-helical conformation. DDR2/CD167b autophosphorylation was induced by the collagen X triple-helical region but not the NC1 domain, indicating that the triple-helical region of collagen X contains a specific DDR2 binding site that is capable of receptor activation. DDR2/CD167b is induced during stellate cell activation and implicate the phosphorylated receptor as a mediator of MMP-2 release and growth stimulation in response to type I collagen. Moreover, type I collagen-dependent upregulation of DDR2/CD167b expression establishes a positive feedback loop in activated stellate cells, leading to further proliferation and enhanced invasive activity.

Form:PBS

Molecular Weight:77.1 kDa

Sequences:Arg422-Glu855

Purity:> 95% by HPLC

Concentration:

Endotoxin Level:<1.0 EU per 1 ug of protein (determined by LAL method)

Storage:Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.